



Higher Maths Formula List

Exponential & Logarithmic Functions

$$\log_a x + \log_a y = \log_a(xy)$$

$$\log_a x - \log_a y = \log_a\left(\frac{x}{y}\right)$$

$$\log_a x^n = n \log_a x$$

$$\log_a a = 1$$

$$\log_a 1 = 0$$



Trigonometry

Trigonometric exact values for common angles in degrees.

Angle	0	30°	45°	60°	90°
	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$
<i>Sin</i>	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1
<i>Cos</i>	0	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
<i>Tan</i>	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	<i>No value</i>



The values in the above table are generated from these triangles.



Degrees	30°	45°	60°	90°	120°	150°	180°
Radians	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{5\pi}{6}$	π

Degrees	180°	210°	240°	270°	300°	330°	360°
Radians	π	$\frac{7\pi}{6}$	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	$\frac{11\pi}{6}$	2π

The Addition Formulas

$$\sin(x + a) = \sin x \cos a + \cos x \sin a$$

$$\sin(x - a) = \sin x \cos a - \cos x \sin a$$

$$\cos(x + a) = \cos x \cos a - \sin x \sin a$$

$$\cos(x - a) = \cos x \cos a + \sin x \sin a$$

The Double Angle Formulas

$$\sin 2A = 2\sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A$$

$$\cos 2A = 2\cos^2 A - 1$$

$$\cos 2A = 1 - 2\sin^2 A$$

Trigonometric Identities

$$\sin^2 x + \cos^2 x = 1$$

$$\frac{\sin x}{\cos x} = \tan x$$



Vectors

For the vector $\underline{a} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$, the magnitude of \underline{a} , written $|\underline{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$

For the vectors $\underline{a} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$ and $\underline{b} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}$ the scalar product of \underline{a} & \underline{b} , written $\underline{a} \cdot \underline{b}$, is given

by: $\underline{a} \cdot \underline{b} = a_1b_1 + a_2b_2 + a_3b_3$ and $\underline{a} \cdot \underline{b} = |\underline{a}||\underline{b}|\cos\theta$

where θ is the angle between the positive direction of \underline{a} & \underline{b}

Quadratic Functions

For the quadratic function $ax^2 + bx + c$ the discriminant is defined by $b^2 - 4ac$.

If $b^2 - 4ac < 0$ the quadratic function has no real roots

If $b^2 - 4ac = 0$ the quadratic function has one real root (referred to as equal roots)

If $b^2 - 4ac > 0$ the quadratic function has two real roots

Differentiation

The Chain Rule

To differentiate the function $y = (ax + b)^n$ we use the chain rule. $\frac{dy}{dx} = na(ax + b)^{n-1}$

$f(x)$	$f'(x)$
$\sin ax$	$a \cos ax$
$\cos ax$	$-a \sin ax$



Rates of Change

Let $s(t)$ be a function which gives distance, d , at time t , called displacement.

Then $V(t) = s'(t)$ i.e. velocity is the derivative of displacement

And $A(t) = V'(t)$ i.e. acceleration is the derivative of velocity

Stationary Points

Stationary points occur where $f'(x) = 0$ or $\frac{dy}{dx} = 0$

Stationary points can be a minimum turning point, maximum turning point, or, less commonly, a point of inflection.

Integration

To integrate the function $y = (ax + b)^n$ we use the reverse chain rule.

$$\int (ax + b)^n dx = \frac{(ax + b)^{n+1}}{(n + 1) \cdot a} + c$$

$f(x)$	$\int f(x) dx$
$\sin ax$	$-\frac{1}{a} \cos ax$
$\cos ax$	$\frac{1}{a} \sin ax$



For the curve $y = f(x)$ the area between the curve, the x -axis and the limits a & b is given by:

$$\text{Area} = \int_a^b f(x) dx$$

The area between two curves, $f(x)$ & $g(x)$, is given by:

$$\int_a^b (g(x) - f(x)) dx$$

where $g(x)$ is the upper function and $f(x)$ is the lower function.

The formula for the area between curves can be written as:

$$\text{Area between curves} = \int_a^b (\text{upper function} - \text{lower function}) dx$$

Straight Lines

The equation of a straight line is given by $y = mx + c$, where m is the gradient and c is the y -intercept.

An alternative form of the equation of a straight line is $y - b = m(x - a)$, where m is the gradient and (a, b) is any point on the line.

Recurrence Relations

A general form of a recurrence relation is $u_{n+1} = au_n + b$.

The sequence generated by the recurrence relation $u_{n+1} = au_n + b$ converges to a limit if $-1 < a < 1$ otherwise the sequence diverges i.e. has no limit.

The limit of the sequence generated by $u_{n+1} = au_n + b$ is given by $L = \frac{b}{1-a}$



Circles

The circle with centre (a, b) and radius r is defined by $(x - a)^2 + (y - b)^2 = r^2$


A special case is the circle centre at the origin $(0, 0)$ and radius r which has equation $x^2 + y^2 = r^2$

The general equation of a circle with centre $(-g, -f)$ and radius $\sqrt{g^2 + f^2 - c}$ is $x^2 + y^2 + 2gx + 2fy + c = 0$

Trigonometric Function Graphs

The Sine Function


The graph of the Sine function on the interval $0 \leq x \leq 360^\circ$.





The Cosine Function

The graph of the Cosine function on the interval $0 \leq x \leq 360^\circ$.



The Tangent Function

The graph of the Tangent function on the interval $0 \leq x \leq 360^\circ$.

