



Integration Techniques

Solutions

The Power Rule

Definite Integrals

Integration by U-Substitution

The Chain Rule

Integration by Parts

Integration using Partial Fractions

Integrating the Trigonometric Functions

Improper Integrals

The Power Rule

Q1) a) $\int x^2 dx = \frac{1}{3}x^3 + c$

c) $\int (x^2 + 3) dx = \frac{1}{3}x^3 + 3x + c$

b) $\int 2x^3 dx = \frac{2}{4}x^4 + c = \frac{1}{2}x^4 + c$

d) $\int 2x^{-3} dx = \frac{2}{-2}x^{-2} + c = -x^{-2} + c$

Q2) a) $\int -x^{-3} dx = \frac{1}{2}x^{-2} + c$

c) $\int \frac{x^{-4}}{-2} dx = \int -\frac{1}{2}x^{-4} dx$
 $= \frac{1}{6}x^{-3} + c$

b) $\int -\frac{1}{2}x^3 + 2x dx = -\frac{1}{8}x^4 + x^2 + c$

d) $2x^{-3} + \frac{1}{4}x^3 dx = \frac{2}{-2}x^{-2} + \frac{1}{16}x^4 + c$
 $= -x^{-2} + \frac{1}{16}x^4 + c$

Q3) a) $\int \frac{5}{x^4} dx = \int 5x^{-4} dx = \frac{5}{-3}x^{-3} + c$
 $= -\frac{5}{3}x^{-3} + c$

b) $\int \frac{3}{2x^2} dx = \int \frac{3}{2}x^{-2} dx$
 $= -\frac{3}{2}x^{-1} + c$

c) $\int \sqrt{x} dx = \int x^{\frac{1}{2}} dx = \frac{1}{\frac{3}{2}}x^{\frac{3}{2}} + c$
 $= \frac{2}{3}x^{\frac{3}{2}} + c$

d) $\int 2\sqrt[4]{x} dx = \int 2x^{\frac{1}{4}} dx = \frac{2}{\frac{5}{4}}x^{\frac{5}{4}} + c$
 $= \frac{8}{5}x^{\frac{5}{4}} + c$



Definite Integrals

$$\begin{aligned}\text{Q1) a) } \int_1^3 2x + 4 \, dx &= [x^2 + 4x]_1^3 = (3^2 + 4 \cdot 3) - (1^2 + 4 \cdot 1) \\ &= 21 - 5 = 16\end{aligned}$$

$$\begin{aligned}\text{b) } \int_0^4 6 - 3x \, dx &= \left[6x - \frac{3}{2}x^2\right]_0^4 = \left(6 \cdot 4 - \frac{3}{2} \cdot 4^2\right) - (0) \\ &= 0 - 0 = 0\end{aligned}$$

$$\begin{aligned}\text{c) } \int_{-1}^2 3x^2 + 5x \, dx &= \left[x^3 + \frac{5}{2}x\right]_{-1}^2 = \left(2^3 + \frac{5}{2} \cdot 2\right) - \left((-1)^3 + \frac{5}{2} \cdot -1\right) \\ &= 13 + 3.5 = 16.5\end{aligned}$$

$$\begin{aligned}\text{d) } \int_{-2}^2 2x^3 \, dx &= \left[\frac{1}{2}x^4\right]_{-2}^2 = \left(\frac{1}{2} \cdot 2^4\right) - \left(\frac{1}{2} \cdot (-2)^4\right) \\ &= 8 - 8 = 0\end{aligned}$$

$$\begin{aligned}\text{e) } \int_0^3 2x + 1 \, dx &= [x^2 + x]_0^3 = (3^2 + 3) - (0) \\ &= 12 - 0 = 12\end{aligned}$$

$$\begin{aligned}\text{f) } \int_{-2}^1 5 - x^3 \, dx &= \left[5x - \frac{1}{4}x^4\right]_{-2}^1 = \left(5 \cdot 1 - \frac{1}{4} \cdot 1^4\right) - \left(5 \cdot (-2) - \frac{1}{4} \cdot (-2)^4\right) \\ &= 4.75 + 14 = 18.75\end{aligned}$$

Integration by U-Substitution

$$\text{Q1) a) Let } I = \int x\sqrt{1+x} \, dx$$

$$u = 1 + x, \text{ giving } \frac{du}{dx} = 1$$

Substitute to give

$$I = \int (u - 1)u^{\frac{1}{2}} \, du$$

$$= \int \left(u^{\frac{3}{2}} - u^{\frac{1}{2}}\right) \, du$$

$$= \frac{2u^{\frac{5}{2}}}{5} - \frac{2u^{\frac{3}{2}}}{3} + c$$

$$= \frac{2(1+x)^{\frac{5}{2}}}{5} - \frac{2(1+x)^{\frac{3}{2}}}{3} + c$$



b) Let $I = \int \frac{2}{\sqrt{x}(x-4)} dx$

$$u = \sqrt{x}, \text{ giving } \frac{du}{dx} = \frac{1}{2\sqrt{x}} \text{ or } 2 du = \frac{dx}{\sqrt{x}}$$

Substitute to give

$$\begin{aligned} I &= \int \frac{4}{(u^2-4)} du \\ &= \int \frac{4}{(u-2)(u+2)} du \\ &= \int \left(\frac{1}{u-2} - \frac{1}{u+2} \right) du \\ &= \ln(u-2) - \ln(u+2) + c \\ &= \ln\left(\frac{u-2}{u+2}\right) + c \\ &= \ln\left(\frac{\sqrt{x}-2}{\sqrt{x}+2}\right) + c \end{aligned}$$

c) Let $I = \int \sin^3 x dx$

$$u = \cos x, \text{ giving } \frac{du}{dx} = -\sin x \text{ or } -du = \sin x dx$$

$$\sin^3 x = \sin x(1 - \cos^2 x) \text{ since } \sin^2 x + \cos^2 x = 1$$

$$I = \int \sin x(1 - \cos^2 x) dx$$

Substitute to give

$$\begin{aligned} I &= \int (u^2 - 1) du \\ &= \frac{u^3}{3} - u + c \\ &= \frac{\cos^3 x}{3} - \cos x + c \end{aligned}$$

d) Let $I = \int \sec^4 x dx$

$$u = \tan x, \text{ giving } \frac{du}{dx} = \sec^2 x \text{ or } du = \sec^2 x dx$$

$$\sec^2 x = 1 + \tan^2 x \text{ from trigonometric identities}$$

$$I = \int (1 + \tan^2 x) \sec^2 x dx$$

Substitute to give

$$\begin{aligned} I &= \int (1 + u^2) du \\ &= u + \frac{u^3}{3} + c \\ &= \tan x + \frac{\tan^3 x}{3} + c \end{aligned}$$



The Chain Rule

Q1) a) $\int (x + 4)^2 dx = \frac{1}{3}(x + 4)^3 + c$

d) $\int (x - 4)^7 dx = \frac{1}{8}(x - 4)^8 + c$

b) $\int (2x + 5)^3 dx = \frac{1}{8}(2x + 5)^4 + c$

e) $\int \left(\frac{2x}{3} + 5\right)^2 dx = \frac{1}{2}\left(\frac{2x}{3} + 5\right)^3 + c$

c) $\int (3x - 5)^2 dx = \frac{1}{9}(3x - 5)^3$

f) $\int (3 - 4x)^3 dx = -\frac{1}{16}(3 - 4x)^4 + c$

Q2) a) $\int \frac{1}{(2x - 3)^{-2}} dx = \int (2x - 3)^2 dx$
 $= \frac{1}{6}(2x - 3)^3 + c$

b) $\int \frac{2}{(5 - 3x)^{-3}} dx = \int 2(5 - 3x)^3 dx$
 $= -\frac{1}{6}(5 - 3x)^4 + c$

c) $\int \frac{1}{(3x - 2)^2} dx = \int (3x - 2)^{-2} dx$
 $= -\frac{1}{3}(3x - 2)^{-1} + c$

d) $\int \sqrt{(5x - 4)} dx = \int (5x - 4)^{\frac{1}{2}} dx$
 $= \frac{2}{15}(5x - 4)^{\frac{3}{2}} + c$

Integration by Parts

Q1) a) $\int x \sin x dx$

Let $u = x$, $\frac{du}{dx} = 1$, giving $du = dx$

Let $\frac{dv}{dx} = \sin x$, $v = -\cos x$

$$\begin{aligned}\int x \sin x dx &= -x \cos x - \cos x(1) dx \\ &= -x \cos x + \int \cos x dx \\ &= -x \cos x + \sin x + c\end{aligned}$$

b) $\int x e^x dx$

Let $u = x$, $\frac{du}{dx} = 1$, giving $du = dx$

Let $\frac{dv}{dx} = e^x$, $v = e^x$

$$\int x e^x dx = x e^x - \int e^x dx = x e^x - e^x + c$$



c) $\int x \sec x \tan x \, dx$

Let $u = x$, $\frac{du}{dx} = 1$, giving $du = dx$

Let $\frac{dv}{dx} = \sec x \tan x$, $v = \sec x$

$$\begin{aligned}\int x \sec x \tan x \, dx &= x \sec x - \int \sec x \, dx \\ &= x \sec x - \ln|\sec x + \tan x| + c\end{aligned}$$

d) $\int \frac{\ln x}{x^3} \, dx$

Let $u = \ln x$, $\frac{du}{dx} = \frac{1}{x}$, giving $du = \frac{1}{x} \, dx$

Let $\frac{dv}{dx} = x^{-3}$, $v = \frac{x^{-2}}{-2}$

$$\begin{aligned}\int \frac{\ln x}{x^3} \, dx &= -\frac{1}{2x^2} \ln x - \frac{1}{2x^2} \left(\frac{1}{x}\right) dx \\ &= -\frac{\ln x}{2x^2} + \int \frac{1}{2} x^{-3} \, dx \\ &= -\frac{\ln x}{2x^2} - \frac{1}{4x^2} + c\end{aligned}$$

e) $\int (x^2 + 1) \ln x \, dx$

Let $u = \ln x$, $\frac{du}{dx} = \frac{1}{x}$, giving $du = \frac{1}{x} \, dx$

Let $\frac{dv}{dx} = x^2 + 1$, $v = \frac{x^3}{3} + x$

$$\begin{aligned}\int (x^2 + 1) \ln x \, dx &= \ln x \left(\frac{x^3}{3} + x\right) - \int \left(\frac{x^3}{3} + x\right) \left(\frac{1}{x}\right) dx \\ &= \left(\frac{x^3}{3} + x\right) \ln x - \int \left(\frac{x^2}{3} + 1\right) dx \\ &= \left(\frac{x^3}{3} + x\right) \ln x - \frac{x^3}{9} - x + c\end{aligned}$$

f) Let $I = \int 12x^2(3 + 2x)^5 \, dx$

Let $u = 12x^2$, $\frac{du}{dx} = 24x$, giving $du = 24x \, dx$

Let $\frac{dv}{dx} = (3 + 2x)^5$, $v = \frac{(3 + 2x)^6}{12}$

$$\begin{aligned}I &= \int 12x^2(3 + 2x)^5 \, dx = 12x^2 \left(\frac{(3 + 2x)^6}{12}\right) - \int 24x \frac{(3 + 2x)^6}{12} \, dx \\ &= x^2(3 + 2x)^6 - \int 2x(3 + 2x)^6 \, dx\end{aligned}$$

Using integration by parts again

Let $J = \int 2x(3 + 2x)^6 \, dx$



Let $u = 2x$, $\frac{du}{dx} = 2$, giving $du = 2dx$

Let $\frac{dv}{dx} = (3 + 2x)^6$, $v = \frac{(3 + 2x)^7}{7}$

$$\begin{aligned} I &= \int 2x(3 + 2x)^6 dx = x \frac{(3 + 2x)^7}{7} - \int \frac{(3 + 2x)^7}{7} dx \\ &= x \frac{(3 + 2x)^7}{7} - \frac{(3 + 2x)^8}{112} + c \end{aligned}$$

$$I = x^2(3 + 2x)^6 - x \frac{(3 + 2x)^7}{7} - \frac{(3 + 2x)^8}{112} + c$$

Q2) a) $\int_0^{\ln 2} x e^{2x} dx$

Let $u = x$, $\frac{du}{dx} = 1$, giving $du = dx$

Let $\frac{dv}{dx} = e^{2x}$, $v = \frac{1}{2} e^{2x}$

$$\begin{aligned} \int_0^{\ln 2} x e^{2x} dx &= \left[\frac{1}{2} e^{2x} (x) \right]_0^{\ln 2} - \int \frac{1}{2} e^{2x} dx \\ &= \left(\frac{1}{2} e^{2 \ln 2} \ln 2 \right) - (0) - \left[\frac{1}{4} e^{2x} \right]_0^{\ln 2} \\ &= \frac{4}{2} \ln 2 - \left(\left(\frac{1}{4} e^{2 \ln 2} \right) - \left(\frac{1}{4} e^0 \right) \right) \\ &= 2 \ln 2 - \frac{4}{4} + \frac{1}{4} \\ &= 2 \ln 2 - \frac{3}{4} \end{aligned}$$

b) $\int_0^{\frac{\pi}{2}} x \sin x dx$

Let $u = x$, $\frac{du}{dx} = 1$, giving $du = dx$

Let $\frac{dv}{dx} = \sin x$, $v = -\cos x$

$$\begin{aligned} \int_0^{\frac{\pi}{2}} x \sin x dx &= [-x \cos x]_0^{\frac{\pi}{2}} - \int_0^{\frac{\pi}{2}} (-\cos x) dx \\ &= \left[-\frac{\pi}{2} \cos \frac{\pi}{2} \right] - (0) + \int_0^{\frac{\pi}{2}} (\cos x) dx \\ &= 0 + [\sin x]_0^{\frac{\pi}{2}} \\ &= \sin \frac{\pi}{2} - \sin 0 = 1 \end{aligned}$$



Integration using Partial Fractions

Q1) a) $\frac{3x+5}{(x+1)(x+2)} \equiv \frac{A}{x+1} + \frac{B}{x+2}$

$$3x + 5 \equiv A(x + 2) + B(x + 1)$$

$$\text{Let } x = -1 \Rightarrow 2 = A$$

$$\text{Let } x = -2 \Rightarrow B = 1$$

$$\int \frac{3x+5}{(x+1)(x+2)} dx = \int \left(\frac{2}{x+1} + \frac{1}{x+2} \right) dx$$

$$= 2 \ln|x+1| + \ln|x+2| + c$$

$$= \ln(|x+1|^2) + \ln|x+2| + c$$

$$= \ln|(x+1)^2(x+2)| + c$$

b) $\frac{3x-1}{(2x+1)(x-2)} \equiv \frac{A}{2x+1} + \frac{B}{x-2}$

$$3x - 1 \equiv A(x - 2) + B(2x + 1)$$

$$\text{Let } x = 2 \Rightarrow 5 = 5B \Rightarrow B = 1$$

$$\text{Let } x = -\frac{1}{2} \Rightarrow -\frac{5}{2} = \frac{5}{2}A \Rightarrow A = -1$$

$$\int \frac{3x-1}{(2x+1)(x-2)} dx = \int \left(\frac{-1}{2x+1} + \frac{1}{x-2} \right) dx$$

$$= -\frac{1}{2} \ln|2x+1| + \ln|x-2| + c$$

$$= \ln|(x-2)\sqrt{2x+1}| + c$$

c) $\frac{2x-6}{(x+3)(x-1)} \equiv \frac{A}{x+3} + \frac{B}{x-1}$

$$2x - 6 \equiv A(x - 1) + B(x + 3)$$

$$\text{Let } x = 1 \Rightarrow -4 = 4B \Rightarrow B = -1$$

$$\text{Let } x = -3 \Rightarrow -12 = -4A \Rightarrow A = 3$$

$$\int \frac{2x-6}{(x+3)(x-1)} dx = \int \left(\frac{3}{x+3} - \frac{1}{x-1} \right) dx$$

$$= 3 \ln|x+3| - \ln|x-1| + c = \ln \left| \frac{(x+3)^3}{x-1} \right| + c$$



$$d) \frac{3}{(2+x)(1-x)}$$

$$\frac{3}{(2+x)(1-x)} \equiv \frac{A}{2+x} + \frac{B}{1-x}$$

$$3 \equiv A(1-x) + B(2+x)$$

$$\text{Let } x = 1 \Rightarrow 3 = 3B \Rightarrow B = 1$$

$$\text{Let } x = -2 \Rightarrow 3 = 3A \Rightarrow A = 1$$

$$\int \frac{3}{(2+x)(1-x)} dx = \int \left(\frac{1}{2+x} + \frac{1}{1-x} \right) dx$$

$$= \ln|2+x| - \ln|1-x| + c = \ln \left| \frac{2+x}{1-x} \right| + c$$

Integrating the Trigonometric Functions

$$Q1) a) \int 5 \cos x \, dx = -5 \sin x + c$$

$$b) \int 3 \sin x \, dx = 3 \cos x + c$$

$$c) \int -\frac{1}{3} \cos x \, dx = \frac{1}{3} \sin x + c$$

$$d) \int 2 \cos(2x) \, dx = -\sin 2x + c$$

$$e) \int 6 \sin\left(\frac{3}{2}x\right) dx = 4 \sin\left(\frac{3}{2}x\right) + c$$

$$f) \int -8 \cos(4x) \, dx = 2 \sin(4x) + c$$

Improper Integrals

$$Q1) a) \int_1^{\infty} \frac{1}{x^3} dx$$

$$= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^3} dx$$

$$= \lim_{t \rightarrow \infty} \left[-\frac{1}{2x^2} \right]_1^t$$

$$= \lim_{t \rightarrow \infty} \left(-\frac{1}{2t^2} + \frac{1}{2} \right) = \frac{1}{2}$$

$$= \lim_{t \rightarrow \infty} \left(-\frac{1}{2t^2} \right) + \lim_{t \rightarrow \infty} \left(\frac{1}{2} \right)$$

$$= 0 + \frac{1}{2} = \frac{1}{2}$$

$$b) \int_2^{\infty} x^{-\frac{3}{2}} dx$$

$$= \lim_{t \rightarrow \infty} \int_2^t x^{-\frac{3}{2}} dx$$

$$= \lim_{t \rightarrow \infty} \left[-2x^{-\frac{1}{2}} \right]_2^t$$

$$= \lim_{t \rightarrow \infty} \left(-2t^{-\frac{1}{2}} + \sqrt{2} \right)$$

$$= \lim_{t \rightarrow \infty} \left(\frac{-2}{\sqrt{t}} + \sqrt{2} \right)$$

$$= \lim_{t \rightarrow \infty} \left(\frac{-2}{\sqrt{t}} \right) + \lim_{t \rightarrow \infty} (\sqrt{2})$$

$$= 0 + \sqrt{2} = \sqrt{2}$$



Q2) a) $\int_0^{\infty} e^x dx$

$$= \lim_{t \rightarrow \infty} \int_0^t e^x dx$$

$$= \lim_{t \rightarrow \infty} [e^x]_0^t$$

$$= \lim_{t \rightarrow \infty} (e^t - 1)$$

$$= \lim_{t \rightarrow \infty} (e^t) + \lim_{t \rightarrow \infty} (-1)$$

$e^t \rightarrow \infty$ as $t \rightarrow \infty$, so the integral diverges

b) $\int_1^{\infty} \frac{1}{\sqrt{x}} dx$

$$= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{\sqrt{x}} dx = \lim_{t \rightarrow \infty} \int_1^t x^{-\frac{1}{2}} dx$$

$$= \lim_{t \rightarrow \infty} [2\sqrt{x}]_1^t$$

$$= \lim_{t \rightarrow \infty} (2\sqrt{t} - 2)$$

$$= \lim_{t \rightarrow \infty} (2\sqrt{t}) + \lim_{t \rightarrow \infty} (-2)$$

$\sqrt{t} \rightarrow \infty$ as $t \rightarrow \infty$, so the integral diverges